

APPROXIMATIONS TO PERMUTATION CLASSES

Robert Brignall
Department of Mathematics
University of Bristol
Bristol, BS8 1TW
UK

Shalosh B. Ekhad
Department of Mathematics
Rutgers University
New Brunswick, NJ 08854
USA

Rebecca Smith
Department of Mathematics
SUNY Brockport
Brockport, NY 14420
USA

Vincent Vatter*
Department of Mathematics
Dartmouth College
Hanover, NH 03755
USA

We investigate a new notion of approximately avoiding a permutation: π *almost avoids* β if one can remove a single entry from π to obtain a β -avoiding permutation.

*Partially supported by EPSRC grant GR/S53503/01.

Date: May 14, 2008

AMS 2000 Subject Classification. 05A05, 05A15

1. INTRODUCTION

The permutation π of length n , written in one-line notation as $\pi(1)\pi(2)\cdots\pi(n)$, is said to *contain* the permutation σ if π has a subsequence that is order isomorphic to σ , and each such subsequence is said to be an *occurrence* of σ in π or simply a σ pattern. For example, $\pi = 491867532$ contains $\sigma = 91672$ because of the subsequence $\pi(2)\pi(3)\pi(5)\pi(6)\pi(9) = 51342$. Permutation containment is easily seen to be a partial order on the set of all (finite) permutations, which we simply denote by \leq . If the permutation π fails to contain σ we say that π *avoids* σ .

A downset in this permutation containment order is referred to as a *permutation class*; in other words, if \mathcal{C} is a permutation class, $\pi \in \mathcal{C}$ and $\sigma \leq \pi$, then $\sigma \in \mathcal{C}$. We denote by \mathcal{C}_n the set $\mathcal{C} \cap S_n$ (the permutations of length n in \mathcal{C}) and we refer to $\sum_{n \geq 0} |\mathcal{C}_n| x^n$ as the *generating function* of \mathcal{C} . Given any set of permutations B , the set $\text{Av}(B) = \{\pi : \beta \not\leq \pi \text{ for all } \beta \in B\}$ forms a permutation class and, conversely, for any permutation class \mathcal{C} there is a unique antichain (set of pairwise incomparable elements) B such that $\mathcal{C} = \text{Av}(B)$; we call this antichain the *basis* of \mathcal{C} .

One area in which permutation classes arise is the study of sorting machines. For example, Knuth [5] showed the class $\text{Av}(231)$ consists precisely of those permutations that can be sorted by a stack (a last-in first-out linear sorting machine), while Tarjan [?] observed that the class of permutations that can be sorted by a network consisting of two parallel queues (first-in first-out linear sorting machines) is $\text{Av}(321)$. A classic result in the field of permutation containment is that the number of permutations in $\text{Av}(231)$ and in $\text{Av}(321)$ of length n are both equal to the n th Catalan number (for bijections between the two sets, see the recent survey by Claesson and Kitaev [2]).

Our interest in this paper is with permutations which “almost lie” in a given permutation class, a concept which we formalize as follows: given a permutation class \mathcal{C} and natural number t , we say that the permutation π *t-almost lies in* (or simply *almost lies in* if $t = 1$) \mathcal{C} if one can remove t (or fewer) entries from π to obtain a permutation that lies in \mathcal{C} . We denote the set of permutations which t -almost lie in \mathcal{C} by \mathcal{C}^{+t} ; note that \mathcal{C}^{+t} is a permutation class itself.

Such approximations to permutation classes have a natural interpretation in terms of sorting machines: if \mathcal{C} consists of those permutations which can be sorted by the machine M then the permutations in \mathcal{C}^{+1} are those which can be sorted by M in parallel with a *one-time use buffer*, which we define as a machine which can hold one entry, once in the sorting process. The classes \mathcal{C}^{+t} for $t \geq 2$ then consist of those permutations which can be sorted by M in parallel with t one-time use buffers. Thus our results show that it is approximately 1% more likely that a randomly ordered deck of playing cards can be sorted by a stack in parallel with a one-time use buffer than it can be sorted by a machine consisting of two queues and a one-time buffer in parallel.

Note that our notion of approximating permutation classes differs from a notion introduced by Noonan [7], who studied permutations with at most one copy of 321; this permutation class, which we denote by $\text{Av}(321^{\leq 1})$, is strictly contained in $\text{Av}(321)^{+1}$.




Figure 1: The shape of a Standard Young Tableau obtained from the RS algorithm applied to a permutation in $\text{Av}(321)^{+1}$ that contains at least one 321 pattern

We provide (in the following two sections) the enumeration of $\text{Av}(231)^{+1}$ and $\text{Av}(321)^{+1}$, i.e., the permutations that can be “almost stack-sorted” and “almost sorted by two parallel queues”, and then conclude with a conjecture. By the usual symmetries of permutations, this completes the enumeration of permutations which almost avoid a pattern of length 3.

2. $\text{Av}(321)^{+1}$

To enumerate the permutations in $\text{Av}(321)^{+1}$ we use the Robinson-Schensted (RS) algorithm. While a detailed description of this algorithm can be found in Sagan’s text [8], a few details suffice for our arguments. First recall that the RS algorithm associates to each permutation π of length n a pair, denoted $(P(\pi), Q(\pi))$, of standard Young tableaux (SYT), each with n cells and of the same shape. We denote the shape of $P(\pi)$ by $\text{sh } P(\pi)$, so in the case where π is of length n , $\text{sh } P(\pi) = \text{sh } Q(\pi)$ is a partition, say $\lambda = (\lambda_1, \dots, \lambda_r)$ of n (which we denote by $\lambda \vdash n$). Schensted [9] proved that the length of the longest decreasing subsequence π is equal to the number of rows of $P(\pi)$, and thus if $\text{sh } P(\pi) = \lambda = (\lambda_1, \dots, \lambda_r)$, then the longest decreasing subsequence of π is of length r . We make use of Greene’s extension of this result which requires a bit of notation: the *conjugate* of the partition $\lambda = (\lambda_1, \dots, \lambda_r)$ is the partition $\lambda' = (\lambda'_1, \dots, \lambda'_s)$ where λ'_i is the length of the i th column of λ , i.e., the number of entries of λ which are at least i .

Theorem 1 (Greene [4]). *If $\text{sh } P(\pi) = \lambda$ and $\lambda' = (\lambda'_1, \dots, \lambda'_s)$ then the entries of π can be partitioned into s disjoint decreasing subsequences π_1, \dots, π_s such that the length of π_i is λ'_i for all $1 \leq i \leq s$.*

The first step in our enumeration is to characterize the shapes of SYT that can arise from a permutation which almost avoids 321.

Proposition 2. *The permutation π lies in $\text{Av}(321)^{+1}$ if and only if $\text{sh } P(\pi)$ is of the form (k, ℓ) or $(k, \ell, 1)$ for some integers $\ell \geq k \geq 1$.*

Proof. No permutation in $\text{Av}(321)^{+1}$ can contain a 4321 pattern as then there would be no entry whose removal gives a 321-avoiding permutation. Similarly, no permutation in $\text{Av}(321)^{+1}$ can contain two disjoint occurrences of 321. Translating into SYT this means

that, for all $\pi \in \text{Av}(321)^{+1}$, $\text{sh } P(\pi)$ has at most one column of length 3 and none of any longer length. This implies that $\text{sh } P(\pi)$ is of one of the two forms listed in the statement of the proposition. For the other direction, note that if $\text{sh } P(\pi) = (k, \ell)$ then π avoids 321, while if $\text{sh } P(\pi) = (k, \ell, 1)$ then π contains a subpermutation of length $k + \ell$ which avoids 321, completing the proof. \square

By Proposition 2 and the RS algorithm we now have that

$$|\text{Av}_n(321)^{+1}| = \sum_{\lambda=(k,\ell)\vdash n} (f^\lambda)^2 + \sum_{\lambda=(k,\ell,1)\vdash n} (f^\lambda)^2,$$

where f^λ denotes the number of SYT of shape λ . The first sum is simply the number of 321-avoiding permutations which, as stated in the introduction, is equal to the n th Catalan number, C_n . To evaluate the second sum we use the Hook Length Formula, which states that for $\lambda \vdash n$, f^λ is equal to $n!$ divided by the product of the hooklengths of cells in the Ferrers diagram of λ . In the case of $\lambda = (k, \ell, 1)$, the product of the hooklengths of cells in the top row is

$$(\ell + 2)\ell \cdots (\ell - k + 2)(\ell - k) \cdots 1 = \frac{(\ell + 2)\ell!}{(\ell - k + 1)},$$

the product for the middle row is $(k + 1)(k - 1)!$, and the solitary cell in the bottom row has a hooklength of 1. Thus we have:

$$\begin{aligned} |\text{Av}_n(321)^{+1}| &= C_n + \sum_{(k,\ell,1)\vdash n} \left(\frac{n!(\ell - k + 1)}{(\ell + 2)\ell!(k + 1)(k - 1)!} \right)^2 \\ &= C_n + \sum_{\ell=\lfloor n/2 \rfloor}^{n-2} \left(\frac{n!(2\ell - n + 2)}{(\ell + 2)\ell!(n - \ell)(n - \ell - 2)!} \right)^2. \end{aligned}$$

An empirical calculation in Maple implies that this function likely has the generating function

$$\frac{1 - 8x + 13x^2 + 24x^3 - 48x^4 - (1 - 6x + x^2 + 34x^3 - 26x^4 - 4x^5)\sqrt{1 - 4x}}{2x^2(1 - x)(1 - 4x)^2}.$$

3. $\text{Av}(231)^{+1}$

Our approach to enumerating the class $\text{Av}(231)^{+1}$ differs significantly from the approach used in the previous section and makes use of the following definition: the entry $\pi(i)$ in π is called *essential* if its removal results in a 231-avoiding permutation. For example, the permutation $\pi = 1742653$ contains two essential entries, $\pi(5) = 4$ and $\pi(7) = 3$. (Also note that if $\pi \in \text{Av}(231)$, then by our definition every entry of π is essential.) As a first step, we make the following observation.




Figure 2: A permutation in $\text{Av}(231)^{+1} \setminus \text{Av}(231)$ in which the greatest element is not involved in a copy of 231.

Proposition 3. *An essential entry in the permutation $\pi \in \text{Av}(231)^{+1}$ participates as the minimum entry in either all or none of the occurrences of 231 in π .*

Proof. Suppose, to the contrary, that $\pi \in \text{Av}(231)^{+1}$ contains indices $i < j < k$ such $\pi(i)\pi(j)\pi(k)$ is order isomorphic to 231 and $\pi(k)$ is essential, but that $\pi(k)$ also participates in another 231 pattern as a non-minimal element. Label the minimal element of this later pattern $\pi(\ell)$. Clearly $\pi(i)\pi(j)\pi(\ell)$ is also order isomorphic to 231, so $\pi - \pi(k) \notin \text{Av}(231)$, a contradiction to the assumption that $\pi(k)$ is essential. \square

By Proposition 3 we can divide the essential entries of a permutation $\pi \in \text{Av}(231)^{+1}$ into *small essential entries*, which participate as the minimum entry in all occurrences of 231, and *large essential entries*, which participate as the minimum entry in no occurrences of 231.

Proposition 4. *The generating function for permutations in $\text{Av}(231)^{+1} \setminus \text{Av}(231)$ in which the greatest element is not involved in a copy of 231 is given by $2(f - c)xc$, where f denotes the generating function for $\text{Av}(231)^{+1}$ and c denotes the generating function for the Catalan numbers.*

Proof. The plot of a permutation of the specified form can be divided into the greatest entry and two regions, A and B , as depicted in Figure 2. One (but not both) of the two regions A or B must contain a permutation in $\text{Av}(231)^{+1} \setminus \text{Av}(231)$, while the other must contain a 231-avoiding permutation. This leads to the generating function specified in the proposition. \square

Proposition 5. *The generating function for permutations with an essential greatest (resp., leftmost, rightmost, or least) entry is $x^2c' + xc - c + 1$.*

Proof. The cases are all similar, so we count permutations with an essential greatest entry. To construct such a permutation, one must insert a new greatest entry into a 231-avoider (such permutations have the generating function $x^2c' + xc$) without creating a 231-avoider (these have the generating function $c - 1$). \square




Figure 3: A permutation with a small essential entry in which the greatest entry is involved in at least one copy of 231 but is not essential.

For the following results we need a bit of notation; for a permutation $\pi \in S_n$ and sets $A, B \subseteq [n]$, we write $\pi(A \times B)$ for the permutation which is order isomorphic to the subsequence of π which has indices from A and values in B .

Proposition 6. *The generating function for permutations in $\text{Av}(231)^{+1}$ with an essential small entry in which the greatest entry participates in a copy of 231 but is not essential is $x^4 c'^2 + xc(x^2 c' + xc - c + 1)$.*

Proof. Take π of length n specifying the hypotheses, and suppose that $\pi(j) = n$ and that the small essential entry is $\pi(k)$. As n must be involved in at least one copy of 231, $\pi([1, j] \times (\pi(k), n))$ must be nonempty; let $\pi(i)$ denote the greatest entry in this region. By this choice of i , $\pi([1, j] \times (\pi(i), n))$ is empty, and because $\pi(k)$ is essential, $\pi((j, n] \times [1, \pi(i)))$ contains only the point $\pi(k)$.

We therefore have three types of entries: the small essential $\pi(k)$, the entries in $\pi([1, j] \times [1, \pi(i)])$, and the entries in $\pi((j, n] \times (\pi(i), n))$. We further divide these latter regions into A, B, C and D , as indicated in Figure 3.

As the entry n is not essential, one of two situations must occur:

- (S1) π has a point in C , or
- (S2) $\pi(k)$ forms a copy of 231 with two entries from B .

Conversely, any permutation of this form that satisfies (S1) or (S2) is of the desired form.

First we count the permutations satisfying (S1). In this case the points in $A \cup B$ form a 231-avoiding permutation, as do the points in $C \cup D$; thus both sets of points are counted by c . The generating function for arrangements of $\pi(k)$ among the points in $A \cup B$ is therefore $x^2 c'$, and this is the same as the generating function for arrangements of $\pi(k)$ among the points in $C \cup D$. Multiplying these functions together doublecounts the point $\pi(k)$, but fails to count $\pi(n)$, so the total contribution of the permutations of this type satisfying (S1) is $x^4 c'^2$.

Now we need to count the permutations that satisfy (S2) but not (S1). We know that π does not have a point in C and that the points of π in D avoid 231. Furthermore, $\pi(k)$




Figure 4: A permutation with a large essential entry in which the greatest entry participates in a copy of 231 but is not essential.

and the points in $A \cup B$ form a permutation with an essential rightmost entry, so their generating function is $x^2c' + xc - c + 1$ by Proposition 5. After taking into account the contribution of n , permutations of this type contribute $xc(x^2c' + xc - c + 1)$, proving the proposition. \square

Proposition 7. *The generating function for permutations without a small essential entry, with a large essential entry that is not the greatest entry, and in which the greatest entry participates in at least one copy of 231 is given by $x^2c'(x^2c' + xc - c + 1) + (x^2c' + x^2c - c + x + 1)(c - 1)$.*

Proof. By arguments analogous to the proof of Proposition 6 it can be established that these permutations are of the form depicted in Figure 4 (in this figure the essential large entry is $\pi(i)$). Note that, as indicated by the figure, π must contain a point in region C as otherwise the greatest element of π would not lie in a copy of 231. We divide these permutations into two types:

- (L1) there is a copy of 231 containing $\pi(i)$ and points in $A \cup B$ or
- (L2) there is no such copy of 231.

In case (L1), the points in $A \cup B$ must avoid 231 because $\pi(i)$ is essential, so they are counted by c . Thus x^2c' is the generating function for the number of arrangements of $\pi(i)$ together with the points in $A \cup B$. We then have that $\pi(i)$ is a leftmost essential element of the permutation given by it and the points in $C \cup D$, so Proposition 5 shows that these entries are counted by $x^2c' + xc - c + 1$. Multiplying these functions doublecounts $\pi(i)$ but does not count the greatest entry of π , so the contribution of the (L1) permutations is $x^2c'(x^2c' + xc - c + 1)$.

In case (L2), $\pi(i)$ and the points in $A \cup B$ form a nonempty 231-avoiding permutation, and are thus counted by $c - 1$. The points in $C \cup D$ together with $\pi(i)$ form a permutation with an essential leftmost entry, but we cannot directly apply Proposition 5 because if π had a unique point in C then that point would be a small essential entry, and we do not wish to count such permutations. Thus we subtract the generating function for permutations

with an essential leftmost entry of value 2; it is easily seen that this generating function is $x(c - xc - 1)$, and so the contribution of permutations in case (L2) is $(x^2c' + x^2c - c + x + 1)(c - 1)$, completing the proof. \square

Theorem 8. *The generating function for the permutations that almost avoid 231 is*

$$\frac{1 - 5x - 6x^2 + 45x^3 - 24x^4 - (1 + x - 4x^2 + x^3)(1 - 4x)^{3/2}}{-2x^2(1 - 4x)^{3/2}}.$$

Proof. Letting f denote the generating function for permutations that almost avoid 231, Propositions 4–7 show that

$$f = c + 2xc(f - c) + x^2c' + xc - c + 1 + x^4c'^2 + xc(x^2c' + xc - c + 1) + x^2c'(x^2c' + xc - c + 1) + (x^2c' + x^2c - c + x + 1)(c - 1),$$

from which the desired solution follows. \square

4. OPEN PROBLEMS

It appears, based on our computations, that $|\text{Av}_n(321)^{+1}| < |\text{Av}_n(231)^{+1}|$ for all $n \geq 4$, which begs for a combinatorial explanation:

Problem 9. *Construct a length-preserving injection from $\text{Av}(321)^{+1}$ to $\text{Av}(231)^{+1}$.*

Finally, we conclude with a conjecture about the exact enumeration problem.

Conjecture 10. *For all t , the generating functions for $\text{Av}(231)^{+t}$ and $\text{Av}(321)^{+t}$ are rational in x and $\sqrt{1 - 4x}$.*

We note that there has been similar work done for sets of permutations with at most t copies of these patterns. Bóna [1] proved that the generating function for $\text{Av}(231^{\leq t})$ is rational in x and $\sqrt{1 - 4x}$ (see also Mansour and Vainshtein [6]). For the pattern 321, Noonan [7] enumerated $\text{Av}(321^{\leq 1})$, while Fulmek [3] counted $\text{Av}(321^{\leq 2})$ and conjectured that the generating function for $\text{Av}(321^{\leq t})$ is rational in x and $\sqrt{1 - 4x}$.

Acknowledgements. We would like to thank Mike Atkinson for fruitful discussions.

REFERENCES

- [1] BÓNA, M. The number of permutations with exactly r 132-subsequences is P -recursive in the size! *Adv. in Appl. Math.* 18, 4 (1997), 510–522.
- [2] CLAESSION, A., AND KITAEV, S. Classification of bijections between 321- and 132-avoiding permutations. arXiv:math.0805.1325v1 [math.CO], 2008.

-
- [3] FULMEK, M. Enumeration of permutations containing a prescribed number of occurrences of a pattern of length three. *Adv. in Appl. Math.* 30, 4 (2003), 607–632.
- [4] GREENE, C. An extension of Schensted’s theorem. *Advances in Math.* 14 (1974), 254–265.
- [5] KNUTH, D. E. *The art of computer programming. Volume 3.* Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1973. Sorting and searching, Addison-Wesley Series in Computer Science and Information Processing.
- [6] MANSOUR, T. Counting occurrences of 132 in an even permutation. *Int. J. Math. Math. Sci.*, 25-28 (2004), 1329–1341.
- [7] NOONAN, J. The number of permutations containing exactly one increasing subsequence of length three. *Discrete Math.* 152, 1-3 (1996), 307–313.
- [8] SAGAN, B. E. *The symmetric group*, second ed., vol. 203 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- [9] SCHENSTED, C. Longest increasing and decreasing subsequences. *Canad. J. Math.* 13 (1961), 179–191.