

Characterising when the simple permutations of a class are monotone griddable

Michael Albert*

Aistis Atminas*

Robert Brignall*

Department of Computer Science
University of Otago
Dunedin
New Zealand

School of Mathematics and Statistics
The Open University
Milton Keynes, MK7 6AA
United Kingdom

February 16, 2017

Abstract

We characterise those permutation classes whose simple permutations are monotone griddable. This characterisation is obtained by identifying a set of nine substructures, at least one of which must occur in any simple permutation containing a long sum of 21s.

1 Introduction

A common route to understanding the structure of a permutation class (and hence, e.g. complete its enumeration) is via its simple permutations, as their structure can be considerably easier to characterise than the entire class. Albert, Atkinson, Homberger and Pantone [4] introduced the notion of *deflatibility* to study this phenomenon: that is, the property that the simples in a given permutation class \mathcal{C} actually belong to a proper subclass $\mathcal{D} \subsetneq \mathcal{C}$. See also Vatter's recent survey [13].

One general case of deflatibility is where the set of simple permutations of a class is finite. Such classes are well-quasi-ordered, finitely based, and have algebraic generating functions [1], and via a Ramsey-type result for simple permutations [9], it is decidable when a permutation class has this property [10].

In this paper, we look beyond classes with finitely many simples to those whose simples are 'monotone griddable', and prove the following characterisation. We postpone formal definitions until later, but see Figure 1 for examples of the structures mentioned.

Theorem 1.1. *The simple permutations in a class \mathcal{C} are monotone griddable if and only if \mathcal{C} does not contain the following structures, or their symmetries:*

- *arbitrarily long parallel sawtooth alternations,*
- *arbitrarily long sliced wedge sawtooth alternations,*
- *proper pin sequences with arbitrarily many turns, and*

*Email addresses: malbert@cs.otago.ac.nz, aistis.atminas@gmail.com, rbrignall@gmail.com.




Figure 1: Examples of the permutations characterising the griddability of simples in Theorem 1.1.

- *spiral proper pin sequences with arbitrarily many extensions.*

In general, classes whose simple permutations are monotone griddable do not immediately possess the range of properties that classes with only finitely many simples do. Indeed, few general properties are known even for classes that are themselves wholly monotone griddable, but this has not diminished the efficacy of the following characterisation for the structural understanding and enumeration of many classes (see, for example [3]).

A *sum of k copies of 21* is the permutation $21\ 43\ 65 \cdots (2k)(2k-1)$, written in one line notation. We will often abbreviate this to $\oplus_k 21$. Similarly, a *skew sum of k copies of 12* is the permutation $\ominus_k 12 = (2k-1)(2k) \cdots 34\ 12$.

Theorem 1.2 (Huczynska and Vatter [11]). *A class \mathcal{C} is monotone griddable if and only if it does not admit arbitrarily long sums of 21 or skew sums of 12. That is, for some k neither $\oplus_k 21$ nor $\ominus_k 12$ belong to \mathcal{C} .*

Aside from the structural information that it provides in its own right, the reason that this simple-to-check characterisation has proved so useful is that the classes to which it has been applied typically in fact possess the stronger property of being *geometrically griddable*. Such classes are well-quasi-ordered, finitely based and have rational generating functions [2]. As we have no direct characterisation for a class to be geometrically griddable, the above theorem (which certainly provides a necessary condition) has often provided enough of a ‘hook’ to solve the task at hand.

It is our hope that Theorem 1.1 can provide a similar ‘hook’ to ease the study of classes whose simple permutations are geometrically griddable. Any such class is known to be well-quasi-ordered, finitely based, and strongly algebraic (meaning that it and every subclass have algebraic generating functions), see Albert, Ruškuc and Vatter [6]. Furthermore, every class with growth rate less than $\kappa \approx 2.20557$, is of this form [6]. For instances of practical enumeration tasks that have exploited the geometric griddability of the simple permutations, see Albert, Atkinson and Vatter [5], and




Figure 2: Examples of the two types of sawtooth alternation and an increasing oscillation in the statement of Theorem 1.3.

Pantone [12].

Our characterisation in Theorem 1.1 relies on the following auxiliary result, which guarantees the existence of certain types of structure in simple permutations that contain a long sum of 21s.

Theorem 1.3. *There exists a function $f(n)$ such that every simple permutation that contains a sum of $f(n)$ copies of 21 must contain a parallel or wedge sawtooth alternation of length $3n$ or an increasing oscillation of length n .*

See Figure 2 for examples of the three types of unavoidable structure. Note that wedge sawtooth alternations are not necessarily simple, so the existence of wedge sawtooth alternations in a permutation class do not guarantee that the simple permutations are not monotone griddable, but Theorem 1.3 does nevertheless provide a sufficient condition.

The rest of this paper is organised as follows. In Section 2 we introduce basic notions and define the structures mentioned in the above results. In Section 3 we give the proof of Theorem 1.3, and in Section 4 we complete the proof of Theorem 1.1. In the final section, we discuss some future directions for this work.

2 Preliminaries

For definitions common to the wider study of permutation patterns, we refer the reader to Bevan’s introduction [8]. For a broader background to the study of permutation classes, see Vatter’s excellent survey [13] in the *Handbook of enumerative combinatorics*.

2.1 Geometry, simplicity and gridding

Critical to our work is the ability to visualise permutations and parts of permutations in the plane. The *plot* of a permutation π of length n is the set of coordinate (or *points*) $(i, \pi(i))$ for $i = 1, \dots, n$. In a slight abuse of notation, we will rarely distinguish between a permutation and its plot.

This exposes an important collection of symmetries that are available – specifically the dihedral group generated by reflections in a vertical, horizontal, or diagonal axis. It is to these symmetries we refer when we make an appeal ‘by symmetry’. In particular to prove Theorem 1.1 it suffices to show that if the simple permutations of a class \mathcal{C} contain arbitrarily long sums of 21, then configurations of one of the specified types must occur.

Given points p_1, \dots, p_k in the plane (typically belonging to the plot of a permutation), denote by $\text{rect}(p_1, \dots, p_k)$ the smallest axes-parallel rectangle that contains them. We call $\text{rect}(p_1, \dots, p_k)$ the *rectangular hull* of p_1, \dots, p_k .

Let \mathcal{R} be any axes-parallel rectangle in the plot of a permutation π . The rectangle \mathcal{R} divides the plot of π into nine regions, and we identify the four ‘corners’ as *NE*, *NW*, *SE* and *SW* of π relative to \mathcal{R} , as illustrated in the following diagram.

NW		NE
	\mathcal{R}	
SW		SE

For the rectangle \mathcal{R} itself, denote by $\pi|_{\mathcal{R}}$ the permutation that is order isomorphic to the points of π contained in \mathcal{R} .

Any point (or collection of points) that lies in one of the four unlabelled regions in the above picture is said to *slice* the rectangle \mathcal{R} . Put formally, if $\mathcal{R} = [a, b] \times [c, d]$ is a rectangle, then the point (x, y) *slices* \mathcal{R} *vertically* if $x \in (a, b)$ and $y \notin [c, d]$, and *horizontally* if $x \notin [a, b]$ and $y \in (c, d)$. By extension, we say that a point *slices* a collection of points in the plane if it slices their rectangular hull.

An *interval* of a permutation π is a (nonempty) set of points $\{(i, \pi(i)) : i \in I\}$ for some set of indices I , such that both I and $\pi(I) = \{\pi(i) : i \in I\}$ form contiguous sets of natural numbers.

We can easily identify an interval geometrically by noting that $\text{rect}((i, \pi(i)) : i \in I)$ cannot be sliced, and must contain only points corresponding to indices from I . Equivalently, the (nonempty) set of points of π belonging to an unsliced axis-parallel rectangle \mathcal{R} form an interval.

Trivially, every singleton of π and the whole of π form intervals. If there are no other intervals and $\pi \neq 1$, then π is said to be *simple*.

Given a permutation σ of length n , and permutations π_1, \dots, π_n , the *inflation* of σ by π_1, \dots, π_n is the permutation obtained by replacing each entry $\sigma(i)$ by a sequence of points order isomorphic to π_i and forming an interval. This permutation is commonly denoted by $\sigma[\pi_1, \dots, \pi_n]$.

The reverse process to inflation (i.e., decomposing a permutation into intervals) forms the basis for the *substitution decomposition*, the essence of which is captured in the following result.

Proposition 2.1 (Albert and Atkinson [1]). *Every permutation π is expressible as the inflation of a unique simple permutation σ . Furthermore, if $|\sigma| \geq 4$, then in the expression*

$$\pi = \sigma[\pi_1, \dots, \pi_n],$$

the intervals π_1, \dots, π_n are also unique.

For a permutation which is the inflation of a simple permutation has length 2 (i.e., $\sigma = 12$ or 21), we do not have the same guarantee of uniqueness of the intervals (although there are methods to recover this if needed). If $\pi = 12[\pi_1, \pi_2]$ for some permutations π_1 and π_2 , then we also write $\pi = \pi_1 \oplus \pi_2$, and say that π is *sum decomposable*. Any permutation that is not sum decomposable is *sum indecomposable*. Similarly, if $\pi = 21[\pi_1, \pi_2]$ we write $\pi = \pi_1 \ominus \pi_2$ and say that π is *skew decomposable*, otherwise π is *skew indecomposable*. Finally, the case where π is both sum and skew

indecomposable corresponds to the case in Proposition 2.1 where the unique simple permutation has length at least 4 (as there are no simple permutations of length 3).

For completeness, we now briefly introduce the notion of griddability. However, we do not actually require this definition in our work (the characterisation provided by Theorem 1.2 suffices). A class \mathcal{C} is *monotone griddable* if there exist integers h and v such that for every permutation $\pi \in \mathcal{C}$, we may divide the plot of π into cells using at most h horizontal and v vertical lines, in such a way as the points in each cell form a monotone increasing or decreasing sequence (or the cell is empty).

2.2 Pin sequences

Following [9], a *pin sequence* is a sequence of points p_1, p_2, \dots in the plot of π such that for each $i \geq 3$, p_i slices $\text{rect}(p_1, \dots, p_{i-1})$. Each pin p_i for $i \geq 3$ has a direction – one of *left*, *right*, *up* or *down* – based on its position relative to the rectangle that it slices. By convention, the pins p_1 and p_2 will be regarded as having no direction.

A *proper pin sequence* is one that satisfies two additional conditions:

- *Maximality*: each pin must be maximal in its direction. For example, if p_i is a right pin, then there are no points further to the right of p_i that slice $\text{rect}(p_1, \dots, p_{i-1})$.
- *Separation*: p_{i+1} must *separate* p_i from $\{p_1, \dots, p_{i-1}\}$. In other words, p_{i+1} must lie horizontally or vertically between $\text{rect}(p_1, \dots, p_{i-1})$ and p_i .

As all pin sequences required in the sequel will be proper, for brevity we will sometimes use the term ‘pin sequence’ to mean a proper pin sequence. We now recall some basic properties of (proper) pin sequences.

Proposition 2.2 (Brignall, Huczynska and Vatter [9]). *In a (proper) pin sequence p_1, \dots, p_m ,*

- (a) p_{i+1} cannot lie in the same or opposite direction as p_i (for all $i \geq 3$);
- (b) p_{i+1} does not slice $\text{rect}(p_1, \dots, p_{i-1})$;
- (c) p_i and p_{i+1} are separated either by p_{i-1} or by each of p_1, \dots, p_{i-2} ; and
- (d) one of the sets of points $\{p_1, \dots, p_m\}$, $\{p_1, \dots, p_m\} \setminus \{p_1\}$, or $\{p_1, \dots, p_m\} \setminus \{p_2\}$ is order isomorphic to a simple permutation.

See the lower-left part of Figure 1 for examples of pin sequences.

The following result is critical to what will follow later. A pin sequence p_1, \dots, p_m in a permutation π is said to be *right reaching* if p_m is the rightmost point of π .

Lemma 2.3 (Brignall, Huczynska and Vatter [9]). *For every simple permutation π and pair of points p_1 and p_2 (unless, trivially, p_1 is the right-most point of π), there is a (proper) right-reaching pin sequence beginning with p_1 and p_2 .*


We now introduce some new terminology relating to pins that we will require for our characterisation. Let p_1, \dots, p_m be a pin sequence in a permutation π . We say that a pin p_i *turns* if the direction of p_i is the same as the direction of p_{i-2} . The significance of this concept is in the following observation.

Lemma 2.4. *If a pin sequence p_1, \dots, p_m contains $3(p+q)$ turns, then the permutation corresponding to p_1, \dots, p_m contains $\oplus_p 21$ or $\ominus_q 12$.*

Proof. For a pin sequence p_1, \dots, p_m , let p denote the length of the longest sum of 21s and q the length of the longest skew sum of 12s that the permutation corresponding to the sequence contains. Let $\ell(p_1, \dots, p_m) = p + q$.

We will prove by induction on k the following statement: every pin sequence p_1, \dots, p_m containing $3k$ turns satisfies $\ell(p_1, \dots, p_m) \geq k$. The lemma will follow directly.

The base case $k = 0$ is trivially true, so let p_1, \dots, p_m be a pin sequence with $3(k + 1)$ turns. Let p_j be the latest pin in the sequence which forms a turn, noting that $j \geq 3(k + 1) + 3 \geq 6$. By symmetry, we may assume that p_j is a right pin (and hence so is p_{j-2}) and p_{j-1} an up pin. Note also that p_1, \dots, p_{j-3} contains at least $3(k + 1) - 3 = 3k$ turns, so by induction we know that $\ell(p_1, \dots, p_{j-3}) \geq k$. We now have the following situation:



By inspection, we see that the pair of points p_{j-1}, p_j forms a copy of 21 that is NE of $\text{rect}(p_1, \dots, p_{j-3})$, and so we may add this copy of 21 to the longest sum of 21s that can be found in $\text{rect}(p_1, \dots, p_{j-3})$. Thus we conclude $\ell(p_1, \dots, p_m) \geq \ell(p_1, \dots, p_j) \geq k + 1$. \square

2.3 The permutations in Theorems 1.1 and 1.3

Sawtooth alternations A *sawtooth alternation of length $3n$* is a permutation on $3n$ points that contains a sum of n copies of 21 placed alongside (horizontally or vertically) a monotone sequence of n points, in such a way as each copy of 21 is sliced by a single entry from the monotone sequence – see the first two illustrations in Figure 2.¹

We divide the family of sawtooth alternations into two types: a *parallel sawtooth alternation* is one in which the monotone sequence is increasing, while a *wedge sawtooth alternation* is one in which the monotone sequence is decreasing. See Figure 2 (on page 3).

It is easy to verify that for $n \geq 2$, the parallel sawtooth alternations of length $3n$ are simple. On the other hand, no wedge sawtooth alternation is simple. This fact underpins the extra work that is required in order to get from Theorem 1.3 to Theorem 1.1.

To recover simplicity in wedge sawtooth alternations, consider the sawtooth alternation shown on the left of Figure 2. The leftmost three points of this permutation forms an interval that is order isomorphic to 312. In order to break this (and every other) interval, we form *sliced* wedge sawtooth alternations in one of three ways: pull the ‘1’ of this 312 below all the other points of the monotone sequence (*type 1*), pull the ‘2’ to the right of all other points of the permutation (*type 2*), or replace the ‘1’ with a new maximal element in the permutation (*type 3*). See the top-right portion of Figure 1 (on page 2). For wedge sawtooth alternations that are oriented differently, we make the analogous definitions by appealing to symmetry.

¹Note that for convenience we are only using the term ‘sawtooth alternation’ in respect of sums of 21. By symmetry there are analogous structures that involve skew sums of 12, but we do not need to give these special names.

Proper pin sequences with turns As defined in the previous subsection, a *turn* in a pin sequence is a pin p_i that has the same direction as p_{i-2} . By Lemma 2.4, a pin sequence that contains a lot of turns also contains a long sum of 21s or skew sum of 12s (or both). See the bottom left portion of Figure 1, where the pins that are turns have been marked with hollow points. By Proposition 2.2(d), pin sequences with turns either correspond to simple permutations, or they correspond to permutations for which we may remove one point to recover a simple permutation.

Increasing oscillations An *increasing oscillation* of length n is a permutation on n points formed by a pin sequence that starts from a copy of 21, and then is entirely made up of right and up pins. (That is, for every $i \geq 5$ the pin is a turn.) There are two increasing oscillations of each length, which may be obtained from one another by symmetry. See the rightmost illustration in Figure 2 (on page 3).

Extended spiral pin sequences A pin sequence p_1, \dots, p_m that contains no turns must either follow the repeating pattern of directions ‘left, up, right, down’ or ‘left, down, right, up’. We call both of these pin sequences *spirals*.

Unlike pin sequences with many turns, spiral pin sequences do not contain long sums of 21 or skew sums of 12 (indeed, spiral pin sequences are contained in the class $\text{Av}(3412, 2143)$). To recover long sums of 21, we add points in specific locations that we will call *extensions*. We will consider two types. For ease of explanation, we will assume that p_{i-1} is an up pin and p_i a right pin; all other cases follow by symmetry.

Type 1: An additional point q is a *type 1 extension* of p_i if $\text{rect}(p_i, q)$ is sliced by p_{i-1} and/or p_{i+1} , and by no other points, and the points p_i, q forms a copy of 21.

Type 2: Three additional points q, r, s that are placed relative to p_i form a *type 2 extension* of p_i if:

- (i) s lies either so that p_{i-1}, s forms a copy of 21 and the only pin slicing $\text{rect}(p_{i-1}, s)$ is p_i , or so that s, p_{i+1} forms a copy of 21 and the only pin slicing $\text{rect}(p_{i+1}, s)$ is p_{i+2} ;
- (ii) q and r form a copy of 21 that is SW of p_i , NE of $\text{rect}(p_1, \dots, p_{i-2})$ and is sliced only by s ; and
- (iii) p_{i+1} separates p_i from q and r , and it is the only pin other than s that slices $\text{rect}(p_i, q, r)$.

A *spiral pin sequence with k extensions* is a spiral pin sequence for which there exists k *distinct* pins to which extensions of either type have been added. See Figure 3 for illustrations of the possible extensions to spirals, and the lower right portion of Figure 1 for two examples of a spiral pin sequence with several extensions.

We observe that any spiral pin sequence with k extensions is simple: starting from the fact that a spiral pin sequence is itself simple, the only possible intervals that could be created when the extensions are added can contain at most one point of the original pin sequence. Furthermore, every point belonging to a Type 1 extension is separated from any other point by at least one point belonging to the spiral, which prevents these points from being contained in a proper interval. For a Type 2 extension, the only exception to this is that the two points q and r (forming the copy of 21) are separated by the third point s , but $\text{rect}(q, r, s)$ is sliced by at least two points from the original spiral.

Lemma 2.5. *A spiral pin sequence with $2k$ extensions contains either $\oplus_k 21$ or $\ominus_k 12$.*




Figure 3: Forming extended spirals. For Type 1, any one of the three hollow points may be added. For Type 2, the copy of 21 is added, together with one of the two slicing points.

Proof. For any of the three possible type 1 extensions q to the pin p_i in Figure 3, we see that p_i, q forms a copy of 21 that is NE of $\text{rect}(p_1, \dots, p_{i-2})$. Similarly, either of the type 2 extensions also provides a copy of 21 that is NE of the same rectangle, $\text{rect}(p_1, \dots, p_{i-2})$. Similar arguments apply by symmetry to the other corners NW, SE and SW. Thus, from $2k$ extensions, by symmetry we can find k that contribute a copy of 21. Furthermore, since any such copy of 21 arising from an extension of p_i is NE or SW of $\text{rect}(p_1, \dots, p_{i-2})$, we conclude that this collection of k copies of 21 forms a copy of $\oplus_k 21$, as required. \square

3 Proof of Theorem 1.3

Let π be a permutation, and let \mathcal{R} be an axes-parallel rectangle in the plot of π . A *sliced copy of 21 that spans \mathcal{R}* is a copy of 21 that is NE or SW of \mathcal{R} and sliced by a point that is NW or SE of \mathcal{R} .

For our proof we will need the following observation:

Observation 3.1. *For the permutation $\pi = \oplus_L 21$, the only non-empty intervals are singletons or the points in a sum of 21, i.e. intervals of the form $\{(i, \pi(i)) : i \in [2k - 1, 2l]\}$ for some $1 \leq k \leq l \leq L$.*

Lemma 3.2. *Let π be a sum indecomposable permutation with $|\pi| > 1$. Any line slicing π must slice a copy of 21.*

Proof. If π were sliced by a line not slicing a copy of 21, then π would equal $\pi_1 \oplus \pi_2$ with π_1 the subpermutation of π below (or left of) the line, and π_2 the subpermutation above (or right of) the line. \square

Lemma 3.3. *In a simple permutation σ , for any point p of σ such that the NE corner of σ relative to p contains 21, there exists a copy of 21 in the NE corner of σ that is sliced by a point in the NW or SE corner.*

Proof. Consider $\sigma|_{NE}$, the restriction of σ to the region NE of p . Since it contains a copy of 21, it is either itself a non-trivial sum indecomposable permutation, or it contains a non-trivial sum indecom-




Figure 4: A sequence of m sliced copies of 21 that span nested rectangles in the same direction immediately yields a wedge sawtooth alternation of length $3m$.

possible component. This component must be sliced in σ by some other point, which necessarily lies in the NW or SE corner of σ relative to p , and the result follows by Lemma 3.2. \square

Lemma 3.4. *Let π be a simple permutation, and let L, m be positive integers. If \mathcal{R} is an axes-parallel rectangle in the plot of π which contains a sum of $L(8m^2)^{8m^2}$ copies of 21, then either π contains a sawtooth alternation of length $3m$, or there is a sum of L copies of 21 contained in a simple permutation within $\pi|_{\mathcal{R}}$.*

Proof. Starting with $\mathcal{R}_0 = \mathcal{R}$ and Σ_0 a set of points in \mathcal{R} forming a sum of $L(8m^2)^{8m^2}$ copies of 21, we will construct a sequence of rectangles $\mathcal{R}_0 \supsetneq \mathcal{R}_1 \supsetneq \dots \supsetneq \mathcal{R}_k$ with $k \leq 8m^2$ with the following properties, for $i \geq 1$.

- (i) Each \mathcal{R}_i contains a set of points $\Sigma_i \subset \Sigma_{i-1}$ that forms a sum of at least $L(8m^2)^{8m^2-i}$ copies of 21;
- (ii) the subpermutation $\pi|_{\mathcal{R}_i}$ is an interval inside $\pi|_{\mathcal{R}_{i-1}}$ (and thus by induction is also an interval inside $\pi|_{\mathcal{R}_0}$).
- (iii) In $\pi|_{\mathcal{R}_{i-1}}$, there exists a copy of 21 that is NE or SW of \mathcal{R}_i , and either forms a sliced copy of 21 spanning \mathcal{R}_i , or is sliced by a point outside \mathcal{R}_0 .

Our construction of rectangles will terminate at \mathcal{R}_k for some $k < 8m^2$ if in $\pi|_{\mathcal{R}_k}$ we can find a copy of a simple permutation that contains a sum of (at least) L copies of 21 or a sawtooth alternation of length $3m$.

Otherwise, our construction terminates when $k = 8m^2$, at which point condition (iii) will guarantee that we have a sum of at least $8m^2$ copies of 21 inside \mathcal{R}_0 (one for each rectangle). Each copy of 21 is either sliced by a point outside \mathcal{R}_0 , or it forms a sliced copy of 21 that spans the next rectangle \mathcal{R}_i in the sequence.

If there are $4m^2$ copies of 21 that are sliced by points outside \mathcal{R}_0 , then we can find m^2 pairs that are sliced on the same side of \mathcal{R}_0 , and we can apply Erdős-Szekeres to find a monotone sequence of m points outside \mathcal{R}_0 slicing copies of 21, giving a sawtooth alternation of length $3m$.

On the other hand, if there are $4m^2$ copies of 21 sliced inside \mathcal{R}_0 , then we can find m^2 copies of 21 that are sliced by points in the same way (i.e., one of the 21 lying NE or SW of the next rectangle, with the slicing point being NW or SE). Because the sequence of rectangles $\mathcal{R}_0, \mathcal{R}_1, \dots$ are nested, the slicing points already form a monotone sequence (see Figure 4), which means that we have in fact found a wedge sawtooth alternation of length $3m^2$ (and hence one of length $3m$).




Figure 5: The two scenarios when $\pi|_{\mathcal{R}_i}$ is sum decomposable: (a) at least $4m^2$ nonsingleton sum components; (b) one component (τ) contains a sum of at least $L(8m^2)^{8m^2-(i+1)}$ copies of 21 and another (τ') contains at least one.

Thus, it now suffices to describe the process to construct \mathcal{R}_{i+1} from \mathcal{R}_i to satisfy (i)–(iii) above. Consider the substitution decomposition of $\pi|_{\mathcal{R}_i}$. First, if $\pi|_{\mathcal{R}_i}$ is skew decomposable, then one of the skew indecomposable components must contain all of Σ_i , so we could restrict $\pi|_{\mathcal{R}_i}$ to this single component, with (i)–(iii) still being satisfied. Thus, without loss of generality, we can assume that $\pi|_{\mathcal{R}_i}$ is skew indecomposable.

Next, suppose that $\pi|_{\mathcal{R}_i}$ is sum decomposable. If Σ_i is contained entirely inside a single sum indecomposable component of $\pi|_{\mathcal{R}_i}$, then we may replace \mathcal{R}_i with the rectangular hull of this component, in which case $\pi|_{\mathcal{R}_i}$ is no longer sum decomposable and we have a different case. Otherwise, Σ_i is distributed across at least two of the sum indecomposable components of $\pi|_{\mathcal{R}_i}$, and note that each such component τ intersects Σ_i in a whole number of 21s.

Since π is simple, every nonsingleton sum indecomposable component of $\pi|_{\mathcal{R}_i}$ must be sliced by a point outside \mathcal{R}_i (and hence outside \mathcal{R}_0), and by Lemma 3.2 this slice must also slice a copy of 21 inside the component. Thus, if we have at least $4m^2$ nonsingleton components in $\pi|_{\mathcal{R}_i}$, we can find a sum of $4m^2$ copies of 21 sliced by points outside \mathcal{R}_0 . Since there are four sides of \mathcal{R}_0 , m^2 of these components have their slicing points on the same side of \mathcal{R}_0 , and Erdős-Szekeres then yields a wedge or parallel sawtooth alternation of length $3m$. See Figure 5(a).

Now we may suppose that we have fewer than $4m^2$ nonsingleton sum indecomposable components in $\pi|_{\mathcal{R}_i}$. By the pigeonhole principle, this means that there is some component τ which contains at least $L(8m^2)^{8m^2-i}/(4m^2) > L(8m^2)^{8m^2-(i+1)}$ copies of 21. We also know (by assumption) that there exists some other nonsingleton component τ' , which must be sliced by some point outside \mathcal{R}_0 , see Figure 5(b). We set \mathcal{R}_{i+1} to be the rectangular hull of τ , and any other nonsingleton component τ' provides the copy of 21 sliced by a point outside \mathcal{R}_0 .

We now turn to the case where $\pi|_{\mathcal{R}_i}$ is sum and skew indecomposable. In this case, consider the substitution decomposition of $\pi|_{\mathcal{R}_i}$, and write $\pi|_{\mathcal{R}_i} = \sigma[\tau_1, \dots, \tau_k]$ where $k = |\sigma|$. By Observation 3.1, each interval τ_j may contain 0, 1, or a multiple of 2 points from Σ_i . First, if $\Sigma_i \subset \tau_j$ for some j , then we may replace \mathcal{R}_i with the rectangular hull of τ_j , and consider this instead.

If for some copy of 21 in Σ_i one of the points is a singleton inside some interval τ_j , then there exists




Figure 6: The scenarios when $\pi|_{\mathcal{R}_i}$ is sum and skew indecomposable: (a) $\pi|_{\mathcal{R}_i}$ contains $2L$ intervals intersecting Σ_i in 1 point; (b) At least $4m^2$ intervals intersect Σ_i in two or more points; (c) τ contains many copies of Σ_i and another interval τ' contains at least 1; (d) τ contains many copies of Σ_i , and another copy of Σ_i is split across two other intervals.

another interval that contains the other point as a singleton. Thus, if we can find at least $2L$ intervals of $\pi|_{\mathcal{R}_i}$ each of which intersects Σ_i in exactly one point, then σ contains a sum of at least L copies of 21, see Figure 6(a). On the other hand, if we can find at least $4m^2$ intervals each of which intersects Σ_i in (at least) two points, then each of these intervals must contain a copy of 21 that is sliced by a point outside \mathcal{R}_0 , and by the earlier argument we can find a sawtooth alternation of length $3m$. See Figure 6(b).

We may now suppose that fewer than $2L$ intervals intersect in exactly one point (so at most L copies of 21 for which this occurs), and fewer than $4m^2$ intersect in two or more points. Since Σ_i comprises at least $L(8m^2)^{8m^2-i}/(4m^2)$ copies of 21, there are at least $L \left((8m^2)^{8m^2-i} - 1 \right)$ copies of 21 in which the '2' and '1' of each pair lie in the same interval. By the pigeonhole principle, there exists an interval τ that contains at least $L \left((8m^2)^{8m^2-i} - 1 \right) / (4m^2) > L(8m^2)^{8m^2-(i+1)}$ copies of 21 from Σ_i . We set \mathcal{R}_{i+1} to be the rectangular hull of τ .

Finally, since τ is not the only interval containing copies of 21 from Σ_i , we can now either find another interval τ' that contains a whole number of copies of 21 from Σ_i , or two intervals that together contain a copy of 21 from Σ_i . In the first case (illustrated in Figure 6(c)), the interval τ' must contain a copy of 21 that is sliced by a point outside \mathcal{R}_0 as required. In the second case (illustrated in Figure 6(d)), the copy of 21 must be sliced either outside \mathcal{R}_0 , or by Lemma 3.3 we can find a copy of 21 in the same region of \mathcal{R}_i relative to \mathcal{R}_{i+1} as the original 21 (NE or SW) which is sliced by a point either NW or SE of \mathcal{R}_{i+1} and within \mathcal{R}_i . This gives us the necessary sliced copy of 21 that spans \mathcal{R}_{i+1} for condition (iii).

This completes the possible cases for the substitution decomposition of $\pi|_{\mathcal{R}_i}$, and hence the proof. Note that since π is a finite permutation, the number of times that we may replace the rectangle \mathcal{R}_i with a smaller rectangle (e.g., when $\pi|_{\mathcal{R}_i}$ is skew decomposable) is bounded. \square

Given a simple permutation π , let $\rho(\pi)$ denote the sum of the lengths of the maximal sawtooth alternations in π of each of the eight types depicted in Figure 2. Our proof of Theorem 1.3 will be complete after we have proved the following lemma.

Lemma 3.5. *For every $m, s \in \mathbb{N}$, there is a function $g(m, s)$ such that every simple permutation π that contains a sum of $g(m, s)$ copies of 21 must either contain an increasing oscillation of length m or satisfy $\rho(\pi) \geq 3s$.*

Proof of Theorem 1.3, given Lemma 3.5. We set $f(n) = g(n, 8n)$ where g is the function from Lemma 3.5. Thus, any π that contains a sum of at least $g(n, 8n)$ copies of 21 must contain an increasing oscillation of length n , or have $\rho(\pi) \geq 24n$. Since $\rho(\pi)$ is the sum of the sizes of the eight different maximal sawtooth alternations that can be found in π , one of these sawtooth alternations has length at least $3n$. \square

Proof of Lemma 3.5. First, for $m = 1, 2, 3$, and any $s \in \mathbb{N}$, we can set $g(m, s) = 2$ (since any simple permutation of length at least four contains an increasing oscillation of length 3). Additionally, for any fixed $m \geq 4$, it is not hard to see that we may take $g(m, 1) = 2$ (since any simple permutation of length at least four contains a sawtooth alternation of length three). Thus, we may now assume that $m \geq 4$ and $s > 1$, and we will show that we may take $g(m, s) = (m + 3) \cdot k + 1$, where $k = (8s^2)^{8s^2} g(m, s - 1)$ by induction on s .




Figure 7: The general set-up in the proof of Lemma 3.5. The crosshatched regions must be empty to avoid sliced copies of 21 that span some rectangle \mathcal{S}_i . The pin p_k is shown here so as to slice \mathcal{S}_3 , and the shaded rectangle denotes $\text{rect}(p_1, \dots, p_{k-2})$.

We start with a simple permutation π which contains a sum of at least $(m+3)k+1$ copies of 21, and denote the points in some longest such sum by Σ . We now partition Σ into $m+3$ disjoint rectangles \mathcal{R}_1 through \mathcal{R}_{k+1} where \mathcal{R}_1 is the bounding rectangle for the first copy of 21 in Σ , and for $2 \leq i \leq k+1$ \mathcal{R}_i is the bounding rectangle for the k least copies of 21 in Σ not contained in any previous \mathcal{R}_j .

If any rectangle \mathcal{R}_i contains a sawtooth alternation of length $3s$, then we are done. Therefore, we may assume that no rectangle contains such a sawtooth alternation, and so in each $\pi|_{\mathcal{R}_i}$ we can find a simple permutation σ_i with $g(m, s-1)$ copies of 21 by Lemma 3.4. For $i = 1, \dots, m+3$, let $\mathcal{S}_i = \text{rect}(\sigma_i)$. If any σ_i contained an increasing oscillation of length m , then we are done, thus by the inductive hypothesis we can assume that $\rho(\sigma_i) = 3s-3$ for $i = 1, \dots, m+3$ (since $\rho(\sigma_i) \geq 3s-3$ and if strictly greater we are done.)

The next few paragraphs are best read in conjunction with Figure 7. Let h denote the horizontal line crossing through the '1' of the initial 21, and v the vertical line crossing through the '2'. Note that the bottom-left corner of π below h and to the left of v is increasing, else we would be able to find a longer sum of 21s in π than Σ . For convenience, we will refer to the L-shaped region below h and/or to the left of v as the *outside* region of π , and the rest of π will be *inside*.

Recall that a sliced copy of 21 that spans \mathcal{S}_j is a copy of 21 that is NE or SW of \mathcal{S}_j , sliced by a point that is NW or SE of \mathcal{S}_j . If we can find a sliced copy of 21 that spans \mathcal{S}_j , then we may append it to one of the eight types of sawtooth alternation in \mathcal{S}_j . Since $\rho(\sigma_j) = 3s-3$, this implies that $\rho(\pi) \geq 3s$ and we are done. Consequently, from now on we will assume that there are no sliced copies of 21 spanning any rectangle \mathcal{S}_j .

Under this assumption, since any point p that slices some \mathcal{S}_i must (by Lemma 3.2) slice a copy of 21, all such slicing points must be below and to the left of the top-right corner of \mathcal{S}_{i+1} , and above and to the right of the bottom-left corner of \mathcal{S}_{i-1} (when these rectangles exist). This implies that a number of regions defined by the four boundary lines of each \mathcal{S}_i must be empty, as identified by the




Figure 8: Up to symmetry, the three situations where p_{j+1} is an up pin but p_j and p_{j+1} are not both in the inside region, all give rise to sliced copies of 21 that span \mathcal{S}_1 or \mathcal{S}_{i_j} . The shaded region denotes $\text{rect}(p_1, \dots, p_j)$. The case where p_{j+1} is a right pin is analogous.

crosshatched areas in Figure 7.

Now consider a shortest right-reaching pin sequence starting from the initial 21 of Σ . We will denote this pin sequence by p_1, p_2, \dots, p_n . For any initial segment p_1, \dots, p_j , let i_j denote the least index (if it exists) such that \mathcal{S}_{i_j} is contained in the NE region of $\text{rect}(p_1, \dots, p_j)$. Observe that $i_1 = i_2 = 1$. For any j satisfying $2 \leq j < n$, the pin p_{j+1} slices the rectangular hull $\text{rect}(p_1, \dots, p_j)$ in such a way as to slice a copy of 21. Thus, whenever p_{j+1} is a right pin or an up pin, we can assume that p_{j+1} does not extend beyond \mathcal{S}_{i_j} . From this, we make two conclusions: first, that $i_{j+1} \leq i_j + 1$, and second, that every \mathcal{S}_i must be sliced by some pin. Note that if p_{j+1} is a down or left pin, then $i_{j+1} = i_j$.

In this pin sequence, we identify pin p_k , which is the first pin such that $\mathcal{S}_1 \subset \text{rect}(p_1, \dots, p_k)$. Clearly p_k must be an up pin or a right pin, and we will assume that it is an up pin, the other case being analogous. We claim that $i_k \leq 4$. If not, then some p_ℓ (with $\ell < k$) slices \mathcal{S}_3 . Since p_k is the earliest up pin that extends at least as far as the top of \mathcal{S}_1 , we conclude that p_ℓ must be a right pin, and can be no higher than the top of \mathcal{S}_1 . However, any such pin must then contribute to a sliced copy of 21 that spans \mathcal{S}_2 . Thus $i_k \leq 4$, and note that this bound is tight, as illustrated by the example placement of p_k and p_{k-1} in Figure 7. Note further that, in any case, none of the pins p_{k+2}, \dots, p_n can slice \mathcal{S}_1 .

We now classify more precisely which $j > k$ can satisfy $i_j < i_{j+1}$. By the earlier comments, this can only happen when p_{j+1} is an up or right pin, and then $i_{j+1} \leq i_j + 1$. We claim that $i_j = i_{j+1}$ unless both p_j and p_{j+1} lie in the inside region.

First, suppose p_{j+1} is a right or up pin in the outside region, then it cannot slice \mathcal{S}_{i_j} (else p_{j+1} slices a copy of 21 in \mathcal{S}_{i_j} , and this sliced copy spans \mathcal{S}_1 , see Figure 8(a)), and it cannot extend beyond \mathcal{S}_{i_j} (else we can find a copy of 21 in $\{p_1, \dots, p_j\}$ sliced by p_{j+1} that spans \mathcal{S}_{i_j} , see Figure 8(b)). Thus, we conclude that \mathcal{S}_{i_j} is NE of $\text{rect}(p_1, \dots, p_{j+1})$, i.e. $i_{j+1} = i_j$.

Next, suppose p_{j+1} is a right or up pin in the inside region, but p_j is outside. By definition, p_j cannot slice \mathcal{S}_{i_j} , from which we conclude that p_{j+1} cannot be contained in \mathcal{S}_{i_j} . Furthermore, p_{j+1} cannot slice \mathcal{S}_{i_j} , else we may take a point in \mathcal{S}_{i_j} together with p_j and p_{j+1} , and form a sliced copy of 21 that spans \mathcal{S}_1 , see Figure 8(c). This completes the claim.

We now identify the least index $k' > k + 1$ such that $i_{k'+1} = i_{k'} + 1$, and note that $i_{k'} \leq 6$ (since $i_k \leq 4$) and none of $p_{k'}, \dots, p_n$ slices \mathcal{S}_1 . By the above argument, the sequence of pins $p_{k'-1}, p_{k'}, p_{k'+1}$ must be ‘up-right-up’ or ‘right-up-right’. Now consider $p_{k'+2}$: if it is a down or left pin, then $\{p_{k'}, p_{k'+1}, p_{k'+2}\}$

forms a sliced copy of 21 that spans \mathcal{S}_1 . Thus, $p_{k'+2}$ must also be an up or right pin, and the same argument applies to all subsequent pins. Thus, the sequence of points $p_{k'-1}, p_{k'}, p_{k'+1}, \dots, p_n$ defines an increasing oscillation. Furthermore, since $i_{k'} \leq 6$ and each subsequent pin can slice at most one more rectangle than its immediate predecessor, since there are $m + 3$ rectangles in total, this oscillation contains at least m points, completing the proof. \square

4 Monotone griddability for simple permutations

In this section, we will complete our proof Theorem 1.1. Starting from Theorem 1.3, our main concern is handling wedge sawtooth alternations since they are not simple. Our key result is the following, whose proof will take up the majority of this section.

Proposition 4.1. *For every $m, p, s \in \mathbb{N}$ there exists a number $h(m, p, s) \in \mathbb{N}$ such that whenever a simple permutation contains a wedge sawtooth alternation of length $3h(m, p, s)$, then it contains a split wedge sawtooth alternation of length $3m$, a proper pin sequence with at least p turns, or a spiral pin sequence with at least s extensions.*

Proof of Theorem 1.1, given Proposition 4.1. First, if \mathcal{C} contains arbitrarily long copies of any of the structures listed, then \mathcal{C} contains arbitrarily long simple permutations that themselves contain arbitrarily long sums of 21 or skew sums of 12. Thus, by Theorem 1.2 the simple permutations in \mathcal{C} cannot be monotone griddable.

Conversely, let $\text{Si}(\mathcal{C})$ denote the set of simple permutations in \mathcal{C} , and suppose that the permutations in $\text{Si}(\mathcal{C})$ are not monotone griddable. By Theorem 1.2, the permutation class formed by taking the closure of the set $\text{Si}(\mathcal{C})$ must contain arbitrarily long sums of 21 or skew sums of 12, and hence there must be simple permutations in \mathcal{C} that contain arbitrarily long sums of 21 or skew sums of 12. If \mathcal{C} contains arbitrarily long sums of 21, then by Theorem 1.3 the simple permutations of \mathcal{C} must contain arbitrarily long sawtooth alternations or increasing oscillations. We are done unless $\text{Si}(\mathcal{C})$ contains only long wedge sawtooth alternations, but in this case we may apply Proposition 4.1 to conclude that \mathcal{C} contains arbitrarily long split wedge sawtooth alternations, proper pin sequences with arbitrarily many turns, or spiral proper pin sequences with arbitrarily many extensions. In any case, we conclude that we have found one of the forbidden substructures specified in Theorem 1.1.

A symmetric argument applies in the case when \mathcal{C} contains arbitrarily long skew-sums of 12. \square

Proof of Proposition 4.1. We will show that $h(m, p, s) = 3mp(2s + 1)$ suffices. Suppose we have a simple permutation π that contains a wedge sawtooth alternation ω of size $3mp(2s + 1)$. By symmetry, we can assume that ω comprises a sum of 21s which are split below, oriented $<$. From left to right, denote the triples of points of ω that form the sliced copies of 21 by $T_1, T_2, \dots, T_{3mp(2s+1)}$.

Now, consider a shortest right-reaching pin sequence $p_1 p_2 \dots p_n$ in π starting at the leftmost sliced copy of 21 of ω . For $i \in \{1, 2, \dots, n\}$, let $t(i)$ be the largest index j such that at least one point of T_j slices or is contained in $\text{rect}(p_1, \dots, p_i)$, and let $\Delta(i) = t(i) - t(i - 1)$ for $i > 1$. Note that $t(1) = 0$, $t(2) = 1$, $t(n) = 3mp(2s + 1)$ and $\sum_{i=2}^n \Delta(i) = t(0) + t(n) = 3mp(2s + 1)$. We are interested in the pins for which $\Delta(i) > 0$. For any p_i with $\Delta(i) > 0$, note that this implies there is at least one point from each of $\Delta(i)$ triples T_j that slice $\text{rect}(p_1, \dots, p_i) \setminus \text{rect}(p_1, \dots, p_{i-1})$.




Figure 9: Identifying points of ω that slice $\text{rect}(p_1, \dots, p_i) \setminus \text{rect}(p_1, \dots, p_{i-1})$ when (a) p_i is a down pin, (b) p_i is a right pin, (c) p_i is an up pin. The shading is as follows: extremality of pins (dark grey), shortcuts (light grey) and regions where Type 1 extensions can occur (crosshatched).

First, if there exists i such that $\Delta(i) \geq m$ then the pin p_i slices a copy of 21 in $\text{rect}(p_1, \dots, p_{i-1})$, which together with $m - 1$ sliced copies of 21 from ω , $T_{t(i)+1}, \dots, T_{t(i)+m-1}$, forms a split wedge sawtooth alternation of length $3m$, of type 1 if p_i is a down pin, type 2 if p_i is a right pin, and type 3 if p_i is an up pin. Consequently, we can now assume that $\Delta(i) < m$ for all i . Letting $M = \{i : \Delta(i) > 0\}$ denote the set of indices i for which $\Delta(i)$ is non-zero, we have $|M| \geq 3p(2s + 1)$.

Next, we are done if the pin sequence contains at least p turns, thus we will assume that there are fewer than p turns in total. Since $|M| \geq 3p(2s + 1)$, there exists a turn-free factor of p_1, \dots, p_n containing at least $3(2s + 1)$ distinct indices i for which $\Delta(i) > 0$. Let this factor be p_k, \dots, p_ℓ , which we will assume forms a clockwise spiral pin sequence (i.e. the directions follow the order up, right, down, left).

We will assume that p_k is a down pin, otherwise we may remove at most three pins from the beginning of p_k, \dots, p_ℓ to recover a spiral sequence beginning with a down pin. The cost of doing this is that p_k, \dots, p_ℓ is now only guaranteed to contain at least $3(2s + 1) - 3 = 6s$ distinct indices i with $\Delta(i) > 0$. Irrespective of this, we have $k \geq 3$ (since the first two pins have no direction), which means that p_k extends from a non-trivial $\text{rect}(p_1, \dots, p_{k-1})$, and therefore p_k slices a copy of 21.

Our discussion is now accompanied by Figure 9. In these diagrams, the dark grey regions contain no points because of the maximality of pins, and the light grey regions contain no points because any such point would enable us to take a ‘shortcut’ in the pin sequence, contradicting our choice of a shortest right-reaching pin sequence. There are also a number of crosshatched areas: these denote regions where the existence of a point will contribute a Type 1 extension.

We now consider any pin p_i (with $k \leq i \leq \ell - 4$) for which $\Delta(i) > 0$, with a view to identifying a Type 1 or Type 2 extension in each case. Note that if p_i is a left pin, then $\Delta(i) = 0$ since $\text{rect}(p_1, \dots, p_{i-1})$ already slices or contains the leftmost sliced copy of 21 in ω . Thus the cases that remain are where p_i is an up, right or down pin.

If p_i is a down pin, then there must be at least one point of ω in a crosshatched region (regions A and B in Figure 9(a)), allowing a Type 1 extension to p_i . Similarly, if p_i is a right pin, then we again conclude that we can find a Type 1 extension to p_i , since there must exist a point of ω regions A or B of Figure 9(b).

This leaves the case where p_i is an up pin, illustrated in Figure 9(c). If any one of the crosshatched regions contains a point, then we can find a Type 1 extension for one of p_{i-3} , p_{i+1} or p_{i+2} , so we now assume that these are empty. Fix some triple T_j which contains a point that slices $\text{rect}(p_1, \dots, p_i) \setminus \text{rect}(p_1, \dots, p_{i-1})$. The '1' of this triple must either coincide with p_{i+1} , or lie in one of the regions A , B or C . This implies the same of the '2', and thence the slicing point must equal p_{i+2} , or lie in one of the regions D , E or F .

If both the '2' and the '1' of T_j lie in C and the slicing point lies in F , then we have a Type 2 extension of p_{i+1} and we are done. There are two other cases to consider: either the slicing point is in region F and the '1' is in region B , or the slicing point is below the pin p_{i+3} (i.e. it equals p_{i+2} or lies in regions D or E). In either case, we have that the '2' and the '1' are sliced by a point that lies below p_{i+3} . We can now substitute the pin p_{i+1} with the '1', and the pin p_{i+2} with the slicing point of T_j , and then the 2 is a Type 1 extension of the '1' (acting as a right pin).²

We have now shown how to find an extension whenever we have a pin p_i with $\Delta(i) > 0$. There are at least 6s of these pins, but the above analysis does not guarantee that the 6s extensions are applied to distinct pins, and extremality may have been (temporarily) violated.

To resolve these issues, define a *spiral* to be a set of four contiguous pins that begins with a down pin. We observe that the above analysis shows us that for any pin p_i with $\Delta(i) > 0$, the pin(s) which can be extended (or substituted and extended) all lie in the same spiral as p_i , or the spiral immediately after p_i . Thus, by restricting our collection of pins p_i with $\Delta(i) > 0$ in p_k, \dots, p_{ell} to a subset for which any pair is separated by a complete spiral of pins, we can ensure that the extensions are applied to distinct pins. Note also that this corrects any issues arising from the violation of extremality.

In order to do this, recall that every left pin p_i satisfies $\Delta(i) = 0$. Thus, we may choose every sixth pin from the collection of 6s pins with $\Delta(i) > 0$, leaving us with a set of at least s pins separated by at least seven points. This, in turn, gives us a set of s distinct pins which have extensions of types 1 or 2, and thus we have formed an extended spiral with s extensions.

Finally, while we cannot appeal to symmetry to cover the case where the pin sequence spirals in the opposite direction, the arguments are similar and so we omit the details. \square

5 Concluding remarks

Decision procedure In this paper we have characterised the classes whose simple permutations are monotone griddable. From this, it should be possible to describe a decision procedure to answer the follow algorithmic problem:

Question 5.1. *Given a finitely based permutation class \mathcal{C} , is it decidable whether the simples in \mathcal{C} are monotone griddable?*

The crux of such an algorithm would likely be to extend existing algorithms that handle pin sequences (such as those given in [7] and [10]) to identify turns and (for spiral pin sequences) extensions.

²In making these pin substitutions, we observe that the resulting sequence violates the extremality condition for proper pin sequences, but it still follows the pattern of a spiral pin sequence, with extensions. Furthermore, we find that $\text{rect}(p_1, \dots, p_{i+6})$ is unchanged, so all future pins will be unaffected by this.

Geometric griddability For a class \mathcal{C} whose simple permutations are all *geometrically* griddable, the class itself is contained in the substitution closure $\langle \text{Si}(\mathcal{C}) \rangle$, whence we can conclude that \mathcal{C} has a number of important properties: it is finitely based, well-quasi-ordered, and is enumerated by an algebraic generating function (see Theorems 4.4 and 6.1 of [6]). For this reason, a geometric analogue to Theorem 1.1 is highly desirable.

Question 5.2. *Does there exist a characterisation of classes whose simple permutations are geometrically griddable?*

This would appear to be a difficult question. In particular, there is no known analogue of Theorem 1.2 to characterise when the class itself is geometrically griddable.

References

- [1] ALBERT, M. H., AND ATKINSON, M. D. Simple permutations and pattern restricted permutations. *Discrete Math.* 300, 1-3 (2005), 1–15.
- [2] ALBERT, M. H., ATKINSON, M. D., BOUVEL, M., RUŠKUC, N., AND VATTER, V. Geometric grid classes of permutations. *Trans. Amer. Math. Soc.* 365, 11 (2013), 5859–5881.
- [3] ALBERT, M. H., ATKINSON, M. D., AND BRIGNALL, R. The enumeration of three pattern classes using monotone grid classes. *Electron. J. Combin.* 19(3), 3 (2012), P20, 34 pp. (electronic).
- [4] ALBERT, M. H., ATKINSON, M. D., HOMBERGER, C., AND PANTONE, J. Deflatability of permutation classes. *Australas. J. Combin.* 64 (2016), 252–276.
- [5] ALBERT, M. H., ATKINSON, M. D., AND VATTER, V. Inflations of geometric grid classes: three case studies. *Australas. J. Combin.* 58 (2014), 24–47.
- [6] ALBERT, M. H., RUŠKUC, N., AND VATTER, V. Inflations of geometric grid classes of permutations. *Israel J. Math.* 205, 1 (2015), 73–108.
- [7] BASSINO, F., BOUVEL, M., PIERROT, A., AND ROSSIN, D. An algorithm for deciding the finiteness of the number of simple permutations in permutation classes. *Adv. in Appl. Math.* 64 (2015), 124–200.
- [8] BEVAN, D. I. Permutation patterns: basic definitions and notation. preprint, arXiv:1506.06673, 2015.
- [9] BRIGNALL, R., HUCZYNSKA, S., AND VATTER, V. Decomposing simple permutations, with enumerative consequences. *Combinatorica* 28 (2008), 385–400.
- [10] BRIGNALL, R., RUŠKUC, N., AND VATTER, V. Simple permutations: decidability and unavoidable substructures. *Theoret. Comput. Sci.* 391, 1–2 (2008), 150–163.
- [11] HUCZYNSKA, S., AND VATTER, V. Grid classes and the Fibonacci dichotomy for restricted permutations. *Electron. J. Combin.* 13 (2006), R54, 14 pp.
- [12] PANTONE, J. The enumerations of permutations avoiding 3124 and 4312. To appear in *Annals of Combinatorics*.
- [13] VATTER, V. Permutation classes. In *Handbook of enumerative combinatorics*, Discrete Math. Appl. (Boca Raton). CRC Press, Boca Raton, FL, 2015, pp. 753–833.